

# Middlesex University Research Repository

An open access repository of  
Middlesex University research

<http://eprints.mdx.ac.uk>

Wilson, Laura J. ORCID logoORCID: <https://orcid.org/0000-0002-3744-6132>, Dimitriou, Lygeri ORCID logoORCID: <https://orcid.org/0000-0002-5093-558X>, Hills, Frank ORCID logoORCID: <https://orcid.org/0000-0001-8235-7545>, Gondek, Marcela B. and Cockburn, Emma (2017) The influence of cold water immersion on markers of recovery following resistance exercise. BASES Conference 2017 – Programme and Abstracts (Journal of Sports Sciences, Vol. 35, Iss. sup1, 2017). In: BASES Conference 2017, 28-29 Nov 2017, Nottingham, UK. . ISSN 0264-0414 [Conference or Workshop Item] (doi:10.1080/02640414.2017.1378421)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/23069/>

## Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

[eprints@mdx.ac.uk](mailto:eprints@mdx.ac.uk)

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

### **The influence of cold water immersion on markers of recovery following resistance exercise.**

Wilson, L., Dimitriou, L., Hills, F.A., Gondek, M.B. & Cockburn, E

The use of cold water immersion (CWI) as a recovery intervention is prevalent amongst athletes. Performance of high volume, heavy load resistance exercise is known to result in disturbances of muscle function, perceptual responses and blood borne parameters. Therefore, the aim of this study was to investigate the influence of CWI on markers of recovery following an acute resistance exercise session. With institutional ethics approval, 16 men (mean age:  $24 \pm 4.7$  years; stature:  $2 \pm 0.05$  m; body mass  $85 \pm 13.5$  kg) completed a lower body resistance exercise session (120 repetitions total) comprising back squats, split squats, hip thrusts and Romanian deadlifts, all performed at 80% predicted 1RM. Participants were match paired into either 10 min CWI ( $10 \pm 0.5$  °C) ( $n=8$ ) or a placebo group ( $n=8$ ) based on a ratio of predicted 1RM and lean mass. Perceptions of soreness and training stress, markers of muscle function and efflux of intracellular proteins were assessed before, and at 24, 48 and 72h post exercise. All data were analysed using magnitude based inferences and 90% confidence intervals. The exercise session resulted in increased muscle soreness, disturbances of muscle function and increases in plasma markers of muscle damage. For perceptions of muscle soreness, there was a possibly trivial effect of CWI at all time points compared to placebo. For isometric peak force there was a possibly harmful effect of CWI at 48h post ( $-11.5$ ;  $\pm 19.6\%$ ) (mean;  $\pm$ CL) compared to placebo. For countermovement jump height there was a possibly harmful effect of CWI at 48h ( $-3.80$ ;  $\pm 10.8\%$ ) and 72h ( $-3.90$ ;  $\pm 10.6\%$ ) post exercise compared to placebo. CWI had a possibly harmful effect on the efflux of creatine kinase (CK) at 24h post ( $1.53$ ;  $x/\div 1.881$ ) (mean;  $x/\div$ CL) and a likely harmful effect at both 48h ( $1.78$ ;  $x/\div 1.898$ ) and 72h ( $2.04$ ;  $x/\div 1.836$ ) post compared to placebo. These results suggest that CWI is not effective at attenuating increases in perceptions of muscle soreness after resistance exercise when compared to a placebo intervention. Further, CWI has a detrimental effect on the CK response, as well as the recovery of isometric peak force and vertical jump performance. The implications of these findings should be carefully considered by athletes and practitioners employing CWI as a recovery strategy following heavy load resistance training.

Wilson, L., Dimitriou, L., Hills, F., Gondek, M. & Cockburn, E. (2017). The influence of cold water immersion on markers of recovery following resistance exercise. *Journal of Sports Sciences*, 35(S1), s66.